

Urban street lighting differentially affects community attributes of airborne and ground-dwelling invertebrate assemblages

Martin T. Lockett¹  | Thérésa M. Jones¹  | Mark A. Elgar¹  | Kevin J. Gaston²  |
Marcel E. Visser³  | Gareth R. Hopkins^{1,4} 

¹School of BioSciences, University of Melbourne, Melbourne, Vic., Australia

²Environment & Sustainability Institute, University of Exeter, Penryn, Cornwall, UK

³Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands

⁴Department of Biology, Western Oregon University, Monmouth, OR, USA

Correspondence

Martin T. Lockett
Email: martylockett@gmail.com

Funding information

Australian Research Council, Grant/Award Number: DP150101191

Handling Editor: Mariana Mayer-Pinto

Abstract

1. The introduction of artificial light at night (ALAN) into natural and urbanised landscapes is a known and highly pervasive disruptor of invertebrate communities. However, the effect of variation in intensity and spectra of ALAN on invertebrate communities inhabiting different spatial niches is little understood. Further, the remarkable ability of ALAN to continue to disrupt biodiversity even in chronically illuminated urban landscapes is not often acknowledged.
2. Here, we simultaneously sampled airborne and ground-dwelling invertebrate assemblages under and between urban street lights to explore the effects on community composition and abundance of (a) proximity to decadal (i.e. long-illuminated) nocturnal street lighting and (b) variation in the spectral output of light.
3. The two assemblages responded differently. For airborne invertebrates, night-time abundance doubled, and night-time assemblage composition was significantly different for traps under, compared with between, street lights. These differences in abundance were not affected by street light intensity, and were absent in day samples, suggesting that even weak ALAN may be causing short-term redistribution of nocturnal invertebrates. Further, the abundance (but not composition) effects of ALAN on airborne invertebrates increased when the street lights emitted a higher proportion of short-wavelength light.
4. In contrast, for ground-dwelling invertebrates, we found only marginal effects of proximity and spectrum of lighting on abundance and no effect on assemblage composition. However, more intense street lighting reduced abundance and altered composition at traps both under and between lights.
5. *Synthesis and Applications.* Public lighting managers must consider ALAN impacts on invertebrate communities not only when introducing ALAN to naïve environments, but also when changing lighting in areas that are highly urbanised and exposed to decades of ALAN. Further, lighting proposals and environmental monitoring of invertebrate communities must take into account the effects on both ground-dwelling and airborne assemblages, as these may respond very differently to the presence, intensity and spectrum of ALAN.

KEYWORDS

artificial light at night, community effects, correlated colour temperature, insects, light emitting diode, light pollution, public lighting

1 | INTRODUCTION

The global shift in the intensity and distribution of artificial light at night (ALAN) is one of the most visible and pervasive anthropogenic stressors on the planet (Falchi et al., 2016; Gaston et al., 2015; Hölker et al., 2010; Longcore & Rich, 2004). ALAN is recognised as a significant community-level stressor of invertebrate assemblages and is linked to local- and landscape-scale shifts in abundance and community composition (Davies et al., 2017; Desouhant et al., 2019; Manfrin et al., 2017; Owens & Lewis, 2018). Moreover, the ecological impact of the presence of ALAN appears to extend beyond the nocturnal environment, driving comparable taxa-wide shifts in diurnal communities (Davies et al., 2012; Knop et al., 2017). Most recently, it has been linked to global declines in invertebrate biodiversity (Boyes et al., 2020; Hölker et al., 2010; Owens et al., 2020).

Exploring the effects of ALAN at the community level is necessary for a greater understanding of the broader ecological consequences of light pollution, but is also challenging (Sanders & Gaston, 2018). Invertebrate studies typically sample from a single spatial stratum (Davies et al., 2012; Knop et al., 2017; van Grunsven et al., 2019; Wakefield et al., 2018); rarely are both airborne and ground-dwelling assemblages sampled simultaneously at a given site (but see Manfrin et al., 2017). If distinct assemblages vary in their response to ALAN, this may mask the scale of its impact. For example, airborne invertebrates suffer direct and often proximate exposure to a light source, such as a street light, leading to behavioural responses including disorientation and attraction/repulsion (Longcore & Rich, 2004), or increased mortality through exhaustion, predation, heat or collision (Eisenbeis, 2006). In contrast, ground-dwelling invertebrates (including flying invertebrates at rest) are typically further from a light source. Thus, for this group, disorientation or attraction effects are likely to be muted, and direct light-related mortality is potentially lower. If ground-dwelling individuals are also shaded by the filtering effects of vegetation and/or terrain (Endler, 1993), such effects may be further reduced.

The spectral characteristics of a light are also important. Invertebrate taxa differ in their spectral sensitivities and associated behavioural responses (Briscoe & Chittka, 2001; Davies et al., 2013; Donners et al., 2018) which may in turn promote spectrum-related shifts in the composition of the communities in which they reside (Longcore et al., 2015; Pawson & Bader, 2014; van Grunsven et al., 2019). Lights emitting a high proportion of short wavelengths typically elicit a stronger invertebrate phototactic response (both in terms of the number of individuals attracted to a light source and the number of taxa affected) compared with lights emitting a higher proportion of long wavelengths (Huemer et al., 2010; Pawson & Bader, 2014; van Grunsven et al., 2014; Wakefield et al., 2018). ALAN-associated behavioural shifts may promote spectrum-mediated

differences in the abundance and composition of invertebrate communities (Davies et al., 2017; van Grunsven et al., 2019). However, the technology of lighting (i.e. the type of light source) is not the sole determinant of its spectral characteristics. Even within a single light source, the spectra of individual lights may vary greatly due to age and manufacturing variances (see Appendix S1: Table S1). To broaden the understanding of the effect of ALAN on urban wildlife, including invertebrates, it is therefore critical that the actual spectral outputs of lighting are incorporated along with other (more anthropocentric) parameters such as measures of illuminance (illumination weighted by the spectral sensitivity of the human eye).

Here, we assessed simultaneously the effect of the proximity and spectral characteristics of street lighting on the composition and abundance of ground-dwelling and airborne invertebrate assemblages. Over 3-day periods in 2016 and 2017, we sampled invertebrate assemblages under (illuminated) and between (darker) street lights (adapted from Davies et al., 2012) in three suburban streetscapes in Melbourne, Australia that have been lit by artificial street lighting nocturnally for decades. Despite the long-term presence of street lighting, we predicted that invertebrate assemblage composition and abundance would still differ under and between street lights (Davies et al., 2012), and that such differences may affect both diurnal and nocturnal communities (*sensu* Davies et al., 2012; Davies et al., 2017; Knop et al., 2017). We predicted that the effects of ALAN would increase with lighting intensity and would be more subtle for ground-dwelling compared with airborne invertebrates due to the differing distances from the light (and variation in potential ground cover). Finally, due to the known variation in phototactic responses of invertebrates with respect to different light spectra, and the preponderance of invertebrate taxa with sensitivity to short-wavelength (UV, blue, green) light (Briscoe & Chittka, 2001), we predicted that lights emitting a higher proportion of short wavelengths would elicit the strongest responses (Huemer et al., 2010; Wakefield et al., 2018).

2 | MATERIALS AND METHODS

2.1 | Study locations

We sampled a single row of 10 street lights (mean \pm SE distance between street lights = 58.28 \pm 5.91 m), at each of three suburban locations (Brunswick, Glenroy and Gowanbrae; Appendix S1: Table S1; Figure S1) in Melbourne, Australia. The locations had a comparable 20-year history of 80-watt, UV-shielded, mercury vapour street lighting, but were each in a different phase of a local council roll-out of 18-watt LED lights. Changes from mercury vapour to LED at some street lights, routine bulb replacement and manufacturing variances meant that the intensity and spectral characteristics of lighting

varied substantially within and between sites and years, including between street lights employing the same technology (Table S1; full spectral data in Lockett et al., 2021). Each location was adjacent to a heavily modified minor watercourse with street-side vegetation comprising low herbaceous cover, dominated by exotic grasses, and scattered trees, dominated by native *Eucalyptus* and *Acacia* species (see Appendix S1: Vegetation characteristics).

2.2 | Invertebrate sampling

We sampled airborne and ground-dwelling invertebrates in 2016 (30 March–2 April) and 2017 (4 April–7 April). These trapping periods were almost exactly 12 months apart, shortly after the equinox, towards the middle of the lunar cycle and over a 72-hr period with <1 mm total precipitation.

2.2.1 | Trap type

We used aerial intercept traps (modified from Young, 2005; Figure S2a) to sample airborne invertebrates and standard pitfall traps (modified from Woodcock, 2005; Figure S2b) to sample ground-dwelling invertebrates (see Appendix S1: Trap characteristics). Pitfall traps consisted of a 90-mm diameter plastic pipe set 125 mm into the ground. Both trap types funnelled invertebrates into a 70-ml vial containing 20 ml of diluted soapy (unscented) water.

2.2.2 | Trap placement

We deployed a pair of aerial intercept and pitfall traps beneath each of the 10 street lights (traps 'under lights'; mean \pm SE horizontal distance from luminaire = 2.70 ± 0.14 m) and a second pair of aerial and pitfall traps between this street light and the next nearest street light (traps 'between lights'; horizontal distance from luminaire = $22.35 \text{ m} \pm 0.85 \text{ m}$; see Appendix S1: Trap characteristics). Traps 'under lights' were offset slightly to avoid dead or dying invertebrates falling from the luminaire into the trap. To ensure we were testing only the effects of light (rather than variation in heat), we placed aerial intercept traps 1 m above the ground (several vertical metres from the luminaire) rather than suspended directly under the luminaire. We repeated this deployment in 2016 and 2017, ensuring that traps were reset as precisely as possible between years. One 'between lights' trap was moved 0.95 m due to mulch covering its 2016 position.

2.2.3 | Sampling procedure

We sampled traps in the hour before sunrise and the hour before sunset (following Davies et al., 2012) over a 72-hr period each year, resulting in a total of 720 aerial and 720 pitfall trap samples (2 trap placements \times 2 daily sampling periods \times 3 days \times 2 years \times 10 street lights \times 3

locations), half of which were collected under and half between street lights. At each sampling event, captured individuals were preserved in 70% ethanol. We sampled all locations simultaneously using three coordinated teams. To minimise possible bias related to sampling time within a location, we reversed the order in which vials were collected within a given location every 24 hr. Two street lights at each location were excluded following repeated loss of vials due to damaged traps.

2.3 | Assessment of lighting

At each street light in each year, we took three measures of light: illuminance (lux; SKL300 light meter; Skye Instruments, Llandrindod Wells, UK), correlated colour temperature (CCT; UPRtek[®] MK350 LED Spectrometer; United Power Research Technology Corporation) and photon flux ($\mu\text{mol photons m}^{-2} \text{ s}^{-1}$; Jaz Irradiance spectrometer; Ocean Optics). Readings were taken at 1 m above the ground directly under the luminaire, on a cloudless night, after astronomical twilight and prior to moonrise. Illuminance was also measured 1 m above the ground at all aerial traps and 0.1 m above the ground at all pitfall traps to confirm that traps under and between lights differed substantially in light levels. We used illuminance and CCT as, while they are calibrated to the human visual system and thus less appropriate when assessing other animal species (Endler, 1990), they are commonly used in lighting industry standards (see e.g. International Electrotechnical Commission, 2014; VicRoads, 2016; Standards Australia, 2018). To provide an unweighted measure of the intensity and spectral output of each light, we measured the total photons emitted ('photon flux', $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) across the UV and visible spectrum (250 nm–750 nm—'total photon flux'), as well as across four 125-nm bins of light comprising the UV (250–375 nm), blue (375–500 nm), yellow (500–625 nm) and red (625–750 nm) spectra. The photon flux in each bin was then divided by total photon flux and defined as 'proportion UV', 'proportion blue', 'proportion yellow' and 'proportion red' respectively. All light measurements were taken with the instrument sensor facing towards zenith.

2.4 | Identification of invertebrates

Samples were reviewed under dissecting microscopes, with specimens identified to family (82%) or higher taxonomic level (18%) using published taxonomic keys (Appendix S1: Table S2). The effects of daily variation were minimised by aggregating each of the three sampling days and nights to give one 'day' sample and one 'night' sample per trap, per year [following Davies et al., 2012; aggregated sample sizes: $n = 192$ (aerial traps) + 192 (pitfall traps)].

2.5 | Analysis

Analyses were conducted in R version 3.5.0 (R Core Team, 2018; R code in Appendix S3). Collembola ($n = 255,850$) and Acari ($n = 15,676$)

were excluded as they were unlikely to be responding to ALAN conditions: the former mostly moving through the soil profile, and the latter trapped as parasites on larger invertebrates. Separate analyses confirmed that neither Collembola nor Acari abundance responded to variation in ALAN (Appendix S2: Table S3). The structure of the remaining data (numerous taxa, each with relatively few specimens) precluded the analysis of the abundance of individual taxa; accordingly, taxon-level responses to ALAN were explored through compositional analysis. Consistent with our experimental aims, and the order of magnitude difference in individuals collected in pitfall compared with aerial traps, we analysed the two trap types separately.

2.5.1 | Abundance

We assessed variation in abundance using generalised linear mixed effects models (packages `LME4` and `LMERTEST`; Bates et al., 2015; Kuznetsova et al., 2017). Aerial trap data were analysed using GLMMs with an assumed Poisson error distribution; pitfall trap data were over-dispersed (`DHARMA` package; Hartig, 2021) and accordingly analysed in GLMMs with a negative binomial distribution. All abundance statistics are reported on the model (log) scale. As illuminance levels under and between lights were strongly bimodal (Appendix S1: Table S1), we treated this as a categorical variable [trap placement: 'between' (baseline) and 'under']. To assess whether proximity to street lighting affected invertebrate abundance comparably during the day and at night, we assessed variation across all samples (including trap placement, sampling period and trap placement \times sampling period interaction term) as fixed effects in the model. In a second set of models, we restricted analyses to night samples only to first assess whether the intensity and spectral characteristics of individual lights affected abundance (including illuminance, CCT and their interactions with trap placement). We then corroborated this analysis using total photon flux and proportion UV, proportion red and proportion yellow. Proportion blue and red were highly negatively correlated ($r = -0.74$, $p < 0.0001$). Where proportion red had a significant effect, the model was rerun with proportion blue instead. In all models, we added location, year and street light ID as random effects. Initial models comprised all possible combinations of the main effects and their interactions ranked by corrected Akaike information criterion (AICc). We applied model averaging (package `MuMIn`, Bartoń, 2018; see Appendix S1: Model averaging) to calculate effect sizes and a 95% confidence interval for each fixed effect (`confint`, `STATS` package; R Core Team, 2018). Effects where the 95% CI did not straddle zero were considered supported. Supported interactions were assessed using post hoc contrasts or slope estimates (`EMMEANS` package, Lenth, 2019) derived from the top model. Predicted values for plotting were generated from final averaged models using the `predict()` function (R Core Team, 2018).

2.5.2 | Community composition

We explored the composition of ground-dwelling and airborne invertebrate assemblages separately using permutational multivariate

analysis of variance (PERMANOVA: `adonis` function, `VEGAN` package; Oksanen et al., 2019). The main model comprised trap placement, sampling period (day, night) and their interaction, stratified using a combined 'location \times year' blocking factor (Oksanen et al., 2019). Models were refined through sequential removal of terms, where $p > 0.10$, and comparison of model AICc scores. Trap placement was of interest and retained even where not significant. Where we identified an effect of trap placement on composition ($p < 0.05$), we performed permutational ($n = 10,000$) pairwise comparisons of 'under' versus 'between' street light samples to identify the contribution of each taxon to the average between-group Bray–Curtis dissimilarity (`simper` function, `VEGAN` package; Bray & Curtis, 1957; Oksanen et al., 2019). As with the abundance analysis, we then repeated analyses with night samples only to assess whether the intensity and spectral characteristics of individual lights affected composition.

3 | RESULTS

We analysed a total of 1,152 vials which, when aggregated per trap, per 3-day period, resulted in 192 aerial and 192 pitfall trap samples, with equal numbers in day and night traps and between years. Excluding the 255,850 Collembola and 15,676 Acari (see Section 2 above), we collected 2074 specimens representing 119 taxa in aerial traps (2016 = 1,256 specimens and 2017 = 818 specimens), and 13,082 specimens representing 130 taxa in pitfall traps (2016 = 8,294 specimens and 2017 = 4,788 specimens; full dataset in Lockett et al., 2021).

3.1 | Abundance

3.1.1 | All samples

The abundance of invertebrates captured in both aerial and pitfall traps was higher during the day than at night; however, in aerial traps this was dependent on trap placement (Table 1a, Figure 1a). In the night samples, abundance in aerial traps placed under lights was twice that of abundance in aerial traps placed between lights (post hoc contrast of under vs. between: $p < 0.0001$). In contrast, abundance in day samples was comparable regardless of whether the trap was placed under or between lights (post hoc contrast: $p = 0.44$; Table 1a; Figure 1a). Abundance in pitfall traps was consistently higher for day compared with night samples, but there was no effect of trap placement (Table 1b; Figure 1b; mean abundance \pm SE = 109.5 \pm 69.5 for day samples; 26.8 \pm 26.4 for night samples).

3.1.2 | Night samples only

Restricting our analyses to night samples, we found main effects of CCT and trap placement for aerial traps, and an interaction between

TABLE 1 Model estimates and 95% confidence intervals for the effects of sampling period (day/night) and trap placement (under/between street lights) on total invertebrate abundance for (a) aerial traps ($n = 96$ traps in 2016, 94 traps in 2017) and (b) pitfall traps ($n = 96$ traps in 2016, 96 traps in 2017). Effects for which the 95% confidence interval did not straddle zero are in bold

	Model-averaged coefficients (\pm SE)	95% CI (Lower and upper bounds)	
(a) Aerial traps			
Intercept	2.65 \pm 0.15	2.18	3.13
Sampling period	-1.15 \pm 0.08	-1.30	-1.00
Trap placement	0.04 \pm 0.05	-0.06	0.14
Sampling period \times Trap Placement	0.65 \pm 0.10	0.46	0.84
(b) Pitfall traps			
Intercept	4.61 \pm 0.20	4.22	5.00
Sampling period	-1.47 \pm 0.08	-1.62	-1.32




FIGURE 1 The relationship between sampling period (day or night), trap placement (under or between street lights) and invertebrate abundance for (a) aerial and (b) pitfall traps. Points = observed data. Circle and whiskers = mean and 95% confidence interval. Note that data points are jittered to avoid overlap

them (Table 2a; Figure 2a). Abundance in aerial traps was negatively related to CCT between street lights, but CCT had no effect under street lights (post hoc estimated slopes: between = -0.25 ± 0.09 ; under = -0.04 ± 0.07). Our analyses further suggest that airborne invertebrate abundance increased with the proportion of longwave 'red' light (625–750 nm) between, but not under, street lights (interaction of trap placement \times proportion red; post hoc estimated slopes: between = 4.44 ± 1.70 ; under = -2.80 ± 1.45 ; Table 2b; Figure 2b). When proportion blue replaced proportion red in the model, we saw the opposite trend (post hoc estimated slopes: between = -4.45 ± 1.70 ; under = 2.80 ± 1.45 ; Appendix S2: Table S4).

Abundance in pitfall traps decreased with increasing total photon flux of the nearest street light; this effect was observed regardless of trap placement (no interaction between total photon flux and trap placement; Table 2b; Figure 2c). Abundance at pitfall traps between

(but not under) street lights was also negatively related to the proportion of UV light (250–375 nm) emitted by the nearest street light (interaction of trap placement \times proportion UV; post hoc estimated slopes: between = -13.66 ± 4.16 ; under = 2.84 ± 3.95 ; Table 2b; Figure 2d). These effects on abundance in pitfall traps were not observed when modelled using illuminance and CCT (Table 2a).

3.2 | Community composition

The composition of day and night samples was significantly different for both aerial and pitfall traps. Sampling period explained approximately 11% of the variance in composition in aerial trap samples, and 16% of variance in pitfall trap samples (Table 3). There was also an effect of trap placement on composition in aerial traps, but this

TABLE 2 Model estimates and 95% confidence intervals for the effects of (a) illuminance, correlated colour temperature (CCT) and their interactions with trap placement; and (b) total photon flux, photon flux proportions (UV, red, yellow) and their interactions with trap placement, on night-time abundance for aerial ($n = 48$ traps in 2016 and 48 traps in 2017) and pitfall traps ($n = 48$ traps in 2016 and 48 traps in 2017). Effects for which the 95% confidence interval did not straddle zero are in bold

(a) Trap placement × Illuminance, CCT			
Aerial traps			
Intercept	1.19 ± 0.39	0.41	1.97
Trap placement	0.68 ± 0.08	0.51	0.84
CCT	-0.22 ± 0.10	-0.43	-0.02
Illuminance (lux)	0.01 ± 0.01	-0.01	0.04
Trap Placement × CCT	0.25 ± 0.10	0.06	0.45
Pitfall traps			
Intercept	3.96 ± 0.21	3.42	4.38
Illuminance (lux)	-0.05 ± 0.01	-0.08	-0.03
(b) Trap placement × total photon flux, photon flux proportions			
Aerial traps			
Intercept	0.54 ± 0.41	-0.28	1.36
Trap placement	1.89 ± 0.32	1.26	2.52
Proportion red	4.98 ± 1.42	2.16	7.80
Proportion UV	-3.92 ± 3.77	-11.4	3.52
Trap placement × Prop. Red	-7.01 ± 1.63	-10.3	-3.77
Trap placement × Prop. UV	-6.82 ± 5.29	-3.59	17.2
Pitfall traps			
Intercept	5.91 ± 1.41	3.11	8.70
Trap placement	-3.84 ± 1.98	-7.74	0.07
Total photon flux	-2.75 ± 0.60	-3.95	-1.55
Proportion UV	-13.6 ± 3.99	-21.5	-5.67
Proportion yellow	-3.43 ± 2.09	-7.56	0.71
Trap placement × Prop. UV	16.3 ± 5.50	5.44	27.2
Trap placement × Prop. yellow	5.73 ± 2.98	-0.14	11.6

explained <1% of the variance across samples (Table 3). There was no comparable effect of trap placement for pitfall traps and no significant sampling period × trap placement interaction regardless of trap type.

Post hoc analyses suggest variation in the composition of airborne samples captured under and between street lights was driven strongly by Diptera, Hemiptera, Hymenoptera and Coleoptera (contribution to dissimilarity index: Diptera families = 0.25; Hemiptera families = 0.19; Hymenoptera families = 0.12; Coleoptera families = 0.10; Appendix S2: Table S5).

In night samples, there was a significant difference in composition between aerial traps placed under and between street lights (Table 4), which explained approximately 2.3% of the variation between samples.

The effects of lighting intensity and spectrum were inconsistent and depended on which measures were used (compare Table 4a and b). In pitfall traps, composition was not significantly affected by trap placement; however, the intensity of the nearest street light (whether measured as illuminance or total photon flux) explained approximately 4.5% of the variation in composition (Table 4a,b).

The composition of airborne samples captured at night under and between street lights was driven strongly by Diptera, Coleoptera, Hymenoptera and Lepidoptera (contribution to dissimilarity index: Diptera families = 0.30; Coleoptera families = 0.16; Hymenoptera families = 0.10; and Lepidoptera families = 0.06; Appendix S2: Table S6).


4 | DISCUSSION

Our data provide three key insights regarding the impact of ALAN for invertebrate assemblages. First, the data confirm that even in an urban streetscape, with more than a 20-year history of exposure to artificial night lighting, the abundance and composition of invertebrate communities continues to be shaped by the presence, intensity and spectral characteristics of lights. Second, through simultaneous sampling of both ground-dwelling and airborne invertebrate assemblages, we demonstrate that the presence of ALAN affected the two assemblages differently: the nocturnal airborne invertebrate assemblage was characterised by differences in abundance and composition between samples collected under versus between street lights. In contrast, we found little 'under versus between' difference in ground-dwelling invertebrates, but a decline in abundance and variation in composition were both related to brighter street lights. A point to note is that while we used two distinct sampling methods, we do not conclude that the airborne and ground-dwelling invertebrate assemblages are themselves separate communities. Given the significant overlap of taxa in the two assemblages [44% (72 out of 163) of families identified were found in both assemblages], and the significant number of alate individuals in pitfall traps (M.T. Lockett, pers. obs.), it is probable that the two assemblages form a single 'biological' community. Thus, the observed effects of ALAN likely include indirect effects; that is, ALAN impacts on one assemblage may have cascading effects on the community structure of the other. Finally, our data highlight the importance of the spectral composition of lighting—the abundance of nocturnal airborne invertebrates captured under lights was positively related to colour temperature (i.e. 'whiter' lights attracted more invertebrates). Conversely, the abundance of airborne invertebrates captured in the darker traps (placed between lights) was negatively related to colour temperature and positively related to the proportion of red light emitted by the nearest street light.

4.1 | Urban street lights continue to fragment invertebrate communities

There has been remarkably little acknowledgment of the power of ALAN to continue to disrupt biological diversity even after long-term

Aerial traps



Pitfall traps




FIGURE 2 The effects of trap placement (under or between street lights) and street light characteristics on night-time invertebrate abundance. *Aerial traps*: (a) the effect of trap placement and correlated colour temperature of nearest street light, and (b) the effect of trap placement and proportion of 'red' photons emitted by nearest street light. *Pitfall traps*: (c) the effect of intensity (total photon flux) of nearest street light, and (d) the effect of trap placement and proportion of ultraviolet light emitted by nearest street light. Points = observed data; lines = smoothed final averaged model predictions with 95% confidence interval

exposure (i.e. decadal or longer). Our data are broadly consistent with the pivotal pitfall study of Davies et al. (2012) in that we found a clear effect of proximity to street lighting on the abundance and (to a much lesser extent) community composition of an invertebrate assemblage. In contrast to their study, we found only weak effects

of ALAN on invertebrate composition, possibly due to the greater taxonomic complexity of our samples (163 here vs. 60 taxa in Davies et al., 2012). In addition, the ground-dwelling assemblage in our study was not affected by proximity to lighting, but its abundance decreased at all traps, and composition shifted, as lighting intensity

	F statistic	R ²	p
(a) Aerial traps			
Trap placement	$F_{1,189} = 2.01$	0.009	0.011
Sampling period	$F_{1,189} = 22.45$	0.106	<0.0001
Sampling period × trap placement	$F_{1,189} = 1.32$	0.006	0.137
(b) Pitfall traps			
Trap placement	$F_{1,191} = 1.59$	0.007	0.089
Sampling period	$F_{1,191} = 36.60$	0.161	<0.0001
Sampling period × trap placement	$F_{1,191} = 1.49$	0.007	0.112

TABLE 3 Summary of results of permutational multivariate analysis of variance (PERMANOVA) of the effects of sampling period and trap placement on composition of trap samples for (a) aerial traps [$n = 96$ (day) + 96 (night)] and (b) pitfall traps [$n = 96$ (day) + 96 (night)]. Significant effects ($\alpha = 0.05$) are in bold

	F statistic	R ²	p
(a) Trap placement × Illuminance, CCT			
Aerial traps			
Trap placement	$F_{1,89} = 2.29$	0.023	0.004
Illuminance	$F_{1,89} = 3.05$	0.031	0.071
CCT	$F_{1,89} = 1.61$	0.016	0.596
Trap placement × CCT	$F_{1,89} = 2.03$	0.021	0.014
Pitfall traps			
Trap placement	$F_{1,93} = 1.48$	0.014	0.093
Illuminance	$F_{1,93} = 4.47$	0.045	0.001
(b) Trap placement × total photon flux, photon flux proportions			
Aerial traps			
Trap placement	$F_{1,90} = 2.34$	0.023	0.003
Total photon flux	$F_{1,90} = 5.36$	0.054	0.222
Trap placement × Total phot. flux	$F_{1,90} = 2.40$	0.024	0.003
Pitfall traps			
Trap placement	$F_{1,93} = 1.48$	0.015	0.099
Total photon flux	$F_{1,93} = 4.43$	0.045	0.011

TABLE 4 Summary of results of permutational multivariate analysis of variance (PERMANOVA) of the effects of (a) illuminance, correlated colour temperature (CCT) and their interactions with trap placement; and (b) total photon flux, photon flux proportions (UV, red, yellow) and their interactions with trap placement, on composition of night-time aerial ($n = 48$ traps in 2016 and 46 traps in 2017) and pitfall trap samples ($n = 48$ traps in 2016 and 48 traps in 2017). Significant effects ($\alpha = 0.05$) are in bold

increased. In both studies, differences were linked to a small number of key taxa that included the Hymenoptera and Coleoptera (although differences in the sampled assemblages meant that we also detected significant shifts in Diptera, Hemiptera and Lepidoptera, while Davies et al. (2012) reported shifts in Opiliones, Isopoda and Amphipoda). An important question is whether these differences are comparable to, or different from, the initial shifts in community structure following introduction of lights to the environment and are thus sustained and permanent changes. A growing body of short-to-medium-term experimental studies suggests analogous changes following exposure to night lighting that appear to persist for multiple generations (Davies et al., 2017; van Grunsven et al., 2019). However, the degree to which these differences are indicative of broadscale fitness declines is largely untested. If street lighting acts as a persistent ecological trap, then it is likely that species that are disproportionately attracted will likely decline over time (Knop et al., 2017; Owens et al., 2020), but trophic interactions may also modify impacts (Bennie et al., 2015, 2018; Perkin et al., 2014). For example, for predators or scavengers, light attraction may provide

trophic benefits that outweigh sublethal ALAN impacts (Willmott et al., 2019).

4.2 | ALAN affects airborne and ground-dwelling assemblages differently

A simultaneous exploration of airborne and ground-dwelling assemblages provides broader insight into the potential fitness impact of street lights. Notably, the assemblages responded differently: nocturnal abundance and overall composition were affected by the proximity of lights for aerial but not for pitfall traps. As hypothesised above, this may have arisen through behavioural differences associated with how these assemblages experience and respond to light. In relatively open streetscapes, strong attraction/avoidance and disorientation behaviours might be expected by airborne invertebrates that have a clear line of sight to a luminaire (Longcore & Rich, 2004). Under this scenario, airborne taxa are drawn towards street lights and away from the darker spaces in between (akin to phototactic

redistribution, *sensu* Eisenbeis, 2006). The absence of this response for ground-dwelling species may be related to the fact that light at pitfall traps was on average 18% less intense than at the aerial traps and thus not as attractive (Appendix S1: Table S1), and the reduced mobility of walking versus flying invertebrates. However, the overall intensity of light from nearby street lights significantly affected pitfall trap abundance: nocturnal ground-dwelling invertebrates in both illuminated (under) and darker (between) spaces were less abundant when the nearest street light was more intense (i.e. a higher total photon flux). This effect was not detected when modelled using illuminance (lux)—a measure weighted to the human spectral sensitivity—potentially underlining the importance of considering all parts of the spectrum when exploring the effects on non-human biota. The composition of nocturnal ground-dwelling invertebrates also shifted with variation in street light intensity, regardless of proximity to the light. The less localised effect of intense light on ground-dwelling invertebrate abundance may indicate more gradual fitness impacts, perhaps through sublethal effects such as reduced fecundity (McLay et al., 2017, 2018) or impaired immune function (Durrant et al., 2015). These effects might disproportionately affect less mobile species or species with reduced geographic ranges and thus could have greater impact on ground-dwelling compared with airborne assemblages.

4.3 | Spectral composition impacts even 'dark' spaces

Apart from behavioural changes associated with the intensity of street lights, our data corroborate the need to consider spectral composition (Davies et al., 2013; Longcore et al., 2015). The abundance of nocturnal airborne invertebrates in the darker traps (placed between lights) was negatively related to colour temperature. Similarly, the abundance of airborne invertebrates at traps between lights was positively related to the proportion of red light emitted by the nearest street light. Both outcomes suggest that shorter wavelength 'white' lights are more efficient at attracting invertebrates away from darker spaces. This result may be attributable to the preponderance of invertebrate taxa with spectral sensitivity peaks in the UV/blue/green part of the spectrum (Briscoe & Chittka, 2001; Davies et al., 2013). The absence of clear spectrum-mediated compositional effects (Table 4a,b) suggests that, rather than partitioning invertebrate assemblages along spectral sensitivity lines, few airborne taxa at our study sites were immune to short-wavelength-mediated phototactic effects. This suggests sensitivity to at least part of the UV/blue/green spectrum is highly conserved, even in those taxa able to exploit longer wavelength light (Davies et al., 2013: their table S1). Thus, notwithstanding the diversity of spectral sensitivities in invertebrate communities, shifting the spectral composition of lights towards longer wavelengths may ameliorate the behavioural impacts of their presence for most species. Given the increasing evidence from across multiple animal taxa (including invertebrates) highlighting the biological relationship between blue light and

physiological processes (Durrant et al., 2015; Grubisic et al., 2019; McLay et al., 2017, 2018), spectrum shifting may assist invertebrates on multiple levels. Disentangling the relative importance of the behavioural and physiological effects of exposure to street lighting in terms of individual fitness is a critical next step to understand the mechanisms underpinning recent evidence for broadscale population declines (Boyes et al., 2020; Knop et al., 2017; Sanders et al., 2020; Senzaki et al., 2020).

4.4 | Management implications

There is an emerging consensus around best practice for minimising the effect of ALAN on wildlife and biodiversity (Gaston et al., 2012; International Dark Skies Association, 2018; Longcore & Rich, 2017). Our study provides further empirical support for these guiding principles, namely that: (a) the presence of ALAN is linked to shifts in community abundance, composition and spatial distribution (*sensu* Sanders & Gaston, 2018); and (b) even moderate amounts of short-wavelength light may impact airborne invertebrate communities (Davies et al., 2017). Our data add further cautionary notes: even in long-illuminated, highly modified urban landscapes, street lighting continues to alter the abundance, spatial distribution and composition of invertebrate communities. Accordingly, public lighting managers and policymakers must take care not only in introducing ALAN to light-naïve environments, but also in changing or augmenting existing public lighting. Further, invertebrate assemblages occupying different spatial niches may respond differently to ALAN: proposed lighting installations and environmental monitoring of invertebrate communities must consider both ground-dwelling and airborne assemblages to capture the full nature and scale of effects. Finally, changing legacy lighting to now-common 4000K LED luminaires will do nothing to ameliorate the behavioural effects of ALAN on airborne invertebrates around street lights; instead it might exacerbate the impact.

ACKNOWLEDGEMENTS

The authors acknowledge the Traditional Owners of the land where this project was carried out, the Wurundjeri people of the Kulin Nations, and pay their respects to their Elders past and present. The thank the City of Moreland and residents of Gowanbrae, Glenroy and Brunswick for hosting this project; Keith Harwood of Ironbark Sustainability for sharing his knowledge of the LED changeover process; and Urban Light Lab volunteers for their assistance with sampling. G.R.H. and this research were funded by an ARC discovery grant (DP150101191) to T.M.J., M.A.E., M.E.V. and K.J.G.

AUTHORS' CONTRIBUTIONS

T.M.J., M.A.E., K.J.G., M.E.V. and G.R.H. conceived the project; T.M.J., G.R.H. and M.T.L. designed the project and carried out data analysis and interpretation of results; Fieldwork and specimen identification was carried out by M.T.L. and G.R.H.; M.T.L., G.R.H. and T.M.J. wrote the manuscript; M.A.E., K.J.G., M.E.V. edited the manuscript.

DATA AVAILABILITY STATEMENT

Data available from the Dryad Digital Repository <https://doi.org/10.5061/dryad.sbccc2fr6s> (Lockett et al., 2021).

ORCID

Martin T. Lockett  <https://orcid.org/0000-0002-7168-4660>
 Thérèse M. Jones  <https://orcid.org/0000-0002-5300-0018>
 Mark A. Elgar  <https://orcid.org/0000-0003-0861-6064>
 Kevin J. Gaston  <https://orcid.org/0000-0002-7235-7928>
 Marcel E. Visser  <https://orcid.org/0000-0002-1456-1939>
 Gareth R. Hopkins  <https://orcid.org/0000-0003-1427-5315>

REFERENCES

- Australia, S. (2018). *Standard SA/SNZ 1158.6:2015 'lighting for roads and public spaces, Part 6: Luminaires – performance'*. SAI Global.
- Bartoń, K. (2018). *MuMIn: Multi-model inference*. R package version 1.40.4. The Comprehensive R Archive Network (CRAN). <https://CRAN.R-project.org/package=MuMIn>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*, 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Bennie, J., Davies, T. W., Cruse, D., Bell, F., & Gaston, K. J. (2018). Artificial light at night alters grassland vegetation species composition and phenology. *Journal of Applied Ecology*, *55*, 442–450. <https://doi.org/10.1111/1365-2664.12927>
- Bennie, J., Davies, T. W., Cruse, D., Inger, R., & Gaston, K. J. (2015). Cascading effects of artificial light at night: Resource-mediated control of herbivores in a grassland ecosystem. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*, 1–9. <https://doi.org/10.1098/rstb.2014.0131>
- Boyes, D. H., Evans, D. M., Fox, R., Parsons, M. S., & Pocock, M. J. O. (2020). Is light pollution driving moth population declines? A review of causal mechanisms across the life cycle. *Insect Conservation and Diversity*, 1–21. <https://doi.org/10.1111/icad.12447>
- Bray, J. R., & Curtis, J. T. (1957). An ordination of upland forest communities of southern Wisconsin. *Ecological Monographs*, *27*, 325–349. <https://doi.org/10.2307/1942268>
- Briscoe, A. D., & Chittka, L. (2001). The evolution of colour vision in insects. *Annual Review of Entomology*, *46*, 471–510. <https://doi.org/10.1146/annurev.ento.46.1.471>
- Davies, T. W., Bennie, J., Cruse, D., Blumgart, D., Inger, R., & Gaston, K. J. (2017). Multiple night-time light-emitting diode lighting strategies impact grassland invertebrate assemblages. *Global Change Biology*, *23*, 2641–2648. <https://doi.org/10.1111/gcb.13615>
- Davies, T. W., Bennie, J., & Gaston, K. J. (2012). Street lighting changes the composition of invertebrate communities. *Biology Letters*, *8*, 764–767. <https://doi.org/10.1098/rsbl.2012.0216>
- Davies, T. W., Bennie, J., Inger, R., Hempel de Ibarra, N., & Gaston, K. J. (2013). Artificial light pollution: Are shifting spectral signatures changing the balance of species interactions? *Global Change Biology*, *19*, 1417–1423. <https://doi.org/10.1111/gcb.12166>
- Desouhant, E., Gomes, E., Mondy, N., & Amat, I. (2019). Mechanistic, ecological, and evolutionary consequences of artificial light at night for insects: Review and prospective. *Entomologia Experimentalis et Applicata*, *167*, 37–58. <https://doi.org/10.1111/eea.12754>
- Donners, M., van Grunsven, R. H. A., Groenendijk, D., van Langevelde, F., Blkker, J. W., Longcore, T., and Veenendaal, E. (2018). Colors of attraction: Modeling insect flight to light behavior. *Journal of Experimental Zoology*, *329*, 434–440. <https://doi.org/10.1002/jez.2188>
- Durrant, J., Michaelides, E. B., Rupasinghe, T., Tull, D., Green, M. P., & Jones, T. M. (2015). Constant illumination reduces circulating melatonin and impairs immune function in the cricket *Teleogryllus commodus*. *PeerJ*, *3*(e1075), 1–23. <https://doi.org/10.7717/peerj.1075>
- Eisenbeis, G. (2006). Artificial night lighting and insects: Attraction of insects to streetlamps in a rural setting in Germany. In C. Rich & T. Longcore (Eds.), *Ecological consequences of artificial night lighting* (pp. 281–304). Island Press.
- Endler, J. A. (1990). On the measurement and classification of colour in studies of animal colour patterns. *Biological Journal of the Linnean Society*, *41*, 315–352. <https://doi.org/10.1111/j.1095-8312.1990.tb00839.x>
- Endler, J. A. (1993). The color of light in forests and its implications. *Ecological Monographs*, *63*, 1–27. <https://doi.org/10.2307/2937121>
- Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C. C. M., Elvidge, C. D., Baugh, K., Portnov, B. A., Rybnikova, N. A., & Furgoni, R. (2016). The new world atlas of artificial night sky brightness. *Science Advances*, *2*, e1600377. <https://doi.org/10.1126/sciadv.1600377>
- Gaston, K. J., Davies, T. W., Bennie, J., & Hopkins, J. (2012). Reducing the ecological consequences of night-time light pollution: Options and developments. *Journal of Applied Ecology*, *49*, 1256–1266. <https://doi.org/10.1111/j.1365-2664.2012.02212.x>
- Gaston, K. J., Visser, M. E., & Hölker, F. (2015). The biological impacts of artificial light at night: The research challenge. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*, 1–6. <https://doi.org/10.1098/rstb.2014.0133>
- Grubisic, M., Haim, A., Bhusal, P., Dominioni, D. M., Gabriel, K. M. A., Jechow, A., Kupprat, F., Lerner, A., Marchant, P., Riley, W., Stebelova, K., van Grunsven, R. H. A., Zeman, M., Zubidat, A. E., & Hölker, F. (2019). Light pollution, circadian photoreception, and melatonin in vertebrates. *Sustainability*, *11*, 1–51. <https://doi.org/10.3390/su11226400>
- Hartig, F. (2021) *DHARMA: Residual diagnostics for hierarchical (multi-level/mixed) regression models, version 0.4.1*. R package. The Comprehensive R Archive Network (CRAN). <https://cran.r-project.org/web/packages/DHARMA/index.html>
- Hölker, F., Wolter, C., Perkin, E. K., & Tockner, K. (2010). Light pollution as a biodiversity threat. *Trends in Ecology & Evolution*, *25*, 681–682. <https://doi.org/10.1016/j.tree.2010.09.007>
- Huemer, P., Kühtreiber, H., & Tarmann, G. (2010). Anlockwirkung moderner Leuchtmittel auf nachtaktive Insekten: Ergebnisse einer Feldstudie in Tirol (Attraction of modern lights on nocturnal insects: Results of a field study in Tyrol). Kooperationsprojekt Tiroler Landesumweltanwaltschaft & Tiroler Landesmuseen Betriebsgesellschaft m.b.H. (Cooperative project between Tirol State Environmental Agency and Tirol State Museum). Innsbruck, Austria. http://www.hellenot.org/fileadmin/user_upload/PDF/WeiterInfo/s/10_AnlockwirkungInsektenFeldstudie_TLMFundLUA.pdf
- International Dark Skies Association. (2018). *Outdoor lighting basics (web-page)*. <http://darksky.org/lighting/lighting-basics/>
- International Electrotechnical Commission. (2014). *Standard IEC 62722-2-1:2014 'Luminaire performance – Part 2-1: Particular requirements for LED luminaires'*. IEC.
- Knop, E., Zoller, L., Ryser, R., Gerpe, C., Hörler, M., & Fontaine, C. (2017). Artificial light at night as a new threat to pollination. *Nature*, *548*, 206–210. <https://doi.org/10.1038/nature23288>
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). ImerTest package: Tests in linear mixed effects models. *Journal of Statistical Software*, *82*, 1–26. <https://doi.org/10.18637/jss.v082.i13>
- Lenth, R. (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.3.5.1. The Comprehensive R Archive Network (CRAN). <https://CRAN.R-project.org/package=emmeans>
- Lockett, M. T., Jones, T. M., Elgar, M. A., Gaston, K. J., Visser, M. E., & Hopkins, G. R. (2021). Data from: Urban street lighting differentially affects community attributes of airborne and ground-dwelling invertebrate assemblages. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.sbccc2fr6s>

- Longcore, T., Aldern, H. L., Eggers, J. F., Flores, S., Franco, L., Hirshfield-Yamanishi, E., Petrinc, L. N., Yan, W. A., & Barroso, A. M. (2015). Tuning the white light spectrum of light emitting diode lamps to reduce attraction of nocturnal arthropods. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370, 20140125. <https://doi.org/10.1098/rstb.2014.0125>
- Longcore, T., & Rich, C. (2004). Ecological light pollution. *Frontiers in Ecology and the Environment*, 2, 191–198. [https://doi.org/10.1890/1540-9295\(2004\)002\[0191:ELP\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2004)002[0191:ELP]2.0.CO;2)
- Longcore, T., & Rich, C. (2017). *Artificial night lighting and protected lands*. National Park Service, U.S. Department of the Interior. <https://irma.nps.gov/DataStore/DownloadFile/582058>
- Manfrin, A., Singer, G., Larsen, S., Weiß, N., van Grunsven, R. H. A., Weiß, N.-S., Wohlfahrt, S., Monaghan, M. T., & Hölker, F. (2017). Artificial light at night affects organism flux across ecosystem boundaries and drives community structure in the recipient ecosystem. *Frontiers in Environmental Science*. <https://doi.org/10.3389/fenvs.2017.00061>
- McLay, L. K., Green, M. P., & Jones, T. M. (2017). Chronic exposure to dim artificial light at night decreases fecundity and adult survival in *Drosophila melanogaster*. *Journal of Insect Physiology*, 100, 15–20. <https://doi.org/10.1016/j.jinsphys.2017.04.009>
- McLay, L. K., Nagarajan-Radha, V., Green, M. P., & Jones, T. M. (2018). Dim artificial light at night affects mating, reproductive output and reactive oxygen species in *Drosophila melanogaster*. *Journal of Experimental Zoology - Part A*, 329, 419–428. <https://doi.org/10.1002/jez.2164>
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2019). *vegan: Community ecology package*. R package version 2.5-6. The Comprehensive R Archive Network (CRAN). <https://CRAN.R-project.org/package=vegan>
- Owens, A. C. S., Cochard, P., Durrant, J., Farnworth, B., Perkin, E. K., & Seymoure, B. (2020). Light pollution is a driver of insect declines. *Biological Conservation*, 241, 108259. <https://doi.org/10.1016/j.biocon.2019.108259>
- Owens, A. C. S., & Lewis, S. M. (2018). The impact of artificial light at night on nocturnal insects: A review and synthesis. *Ecology and Evolution*, 1–22. <https://doi.org/10.1002/ece3.4557>
- Pawson, S. M., & Bader, M.-K.-F. (2014). LED lighting increases the ecological impact of light pollution irrespective of color temperature. *Ecological Applications*, 24, 1561–1568. <https://doi.org/10.1890/14-0468.1>
- Perkin, E. K., Hölker, F., & Tockner, K. (2014). The effects of artificial lighting on adult aquatic and terrestrial insects. *Freshwater Biology*, 59, 368–377. <https://doi.org/10.1111/fwb.12270>
- R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Sanders, D., Frago, E., Kehoe, R., Patterson, C., & Gaston, K. J. (2020). A meta-analysis of biological impacts of artificial light at night. *Nature Ecology & Evolution*. <https://doi.org/10.1038/s41559-020-01322-x>
- Sanders, D., & Gaston, K. J. (2018). How ecological communities respond to artificial light at night. *Journal of Experimental Zoology Part A*, 1–7. <https://doi.org/10.1002/jez.2157>
- Senzaki, M., Barber, J. R., Phillips, J. N., Carter, N. H., Cooper, C. B., Ditmer, M. A., Fristrup, K. M., McClure, C. J. W., Mennitt, D. J., Tyrrell, L. P., Vukomanovic, J., Wilson, A. A., & Francis, C. D. (2020). Sensory pollutants alter bird phenology and fitness across a continent. *Nature*, 587, 605–609. <https://doi.org/10.1038/s41586-020-2903-7>
- van Grunsven, R. H. A., Becker, J., Peter, S., Heller, S., & Hölker, F. (2019). Long-term comparison of attraction of flying insects to streetlights after the transition from traditional light sources to light-emitting diodes in urban and peri-urban settings. *Sustainability*, 11, 1–9. <https://doi.org/10.3390/su11226198>
- van Grunsven, R. H. A., Donners, M., Boekee, K., Tichelaar, I., van Geffen, K. G., Groenendijk, D., Berendse, F., & Veenendaal, E. M. (2014). Spectral composition of light sources and insect phototaxis, with an evaluation of existing spectral response models. *Journal of Insect Conservation*, 1–7. <https://doi.org/10.1007/s10841-014-9633-9>
- VicRoads. (2016). *Specification TCS 065 - 2016 'Specification for the supply of LED road light luminaires'*. VicRoads.
- Wakefield, A., Broyles, M., Stone, E. L., Harris, S., & Jones, G. (2018). Quantifying the attractiveness of broad-spectrum street lights to aerial nocturnal insects. *Journal of Applied Ecology*, 55, 714–722. <https://doi.org/10.1111/1365-2664.13004>
- Willmott, N. J., Henneken, J., Elgar, M. A., & Jones, T. M. (2019). Guiding lights: Foraging responses of juvenile nocturnal orb-web spiders to the presence of artificial light at night. *Ethology*, 125, 289–297. <https://doi.org/10.1111/eth.12852>
- Woodcock, B. A. (2005). Pitfall trapping in ecological studies. In S. R. Leather (Ed.), *Insect sampling in forest ecosystems* (pp. 37–57). Blackwell.
- Young, M. (2005). Insects in flight. In S. R. Leather (Ed.), *Insect sampling in forest ecosystems* (pp. 116–145). Blackwell.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Lockett, M. T., Jones, T. M., Elgar, M. A., Gaston, K. J., Visser, M. E., & Hopkins, G. R. (2021). Urban street lighting differentially affects community attributes of airborne and ground-dwelling invertebrate assemblages. *Journal of Applied Ecology*, 00, 1–11. <https://doi.org/10.1111/1365-2664.13969>